



THE NEW YORK CITY DEPARTMENT OF EDUCATION

Stuyvesant High School

345 Chambers Street, New York, NY 10282-1099
Telephone: (212) 312-4800 Facsimile: (212) 587-3874

April 5, 2024

Dear Parents/Guardians of 11th Grader:

All 12th graders are required to be vaccinated for meningitis meningococcal (menACWY)

Requirements are as follows: one dose only if the 1st dose was received at age 16+ or older. Some students may need a third dose as the first 2 doses were given before the age of 16; proof of immunization is needed and must be signed by a physician or Board of Health person responsible for giving your child these shots. Please see the attached form with the full list of required immunizations.

Records must be brought to Ms. Bullock in Room 371 as soon as possible; failure to provide proof of immunization prior to the start of the school year could result in your child being excluded from school. Ms. Bullock can be reached at (212)317-4800 X3711. Thank you for your prompt attention in this matter.

Brian Moran
Assistant Principal

2023–24: FULL COMPLIANCE

New York State Immunization Requirements for Child Care and School Entrance/Attendance

Notes: For all settings and grades (child care, head start, nursery, 3K, pre-Kindergarten through 12). Intervals between doses of vaccine should be in accordance with the ACIP-recommended immunization schedule for children aged 0 up to 16 years. Doses received more than 4 calendar days before the recommended minimum age or interval are not valid and do not count. This 4-day grace period does not apply to the recommended 28-day minimum interval between doses of live virus vaccines (i.e., MMR, varicella). Refer to the footnotes for dose requirements and specific information about each vaccine. Children enrolling in gradeless classes should meet immunization requirements for their age-equivalent grade. Children who were not in full compliance before the start of the school year must complete requirements according to the ACIP-recommended catch-up schedule in order to remain in child care or school.

VACCINES	CHILD CARE, HEAD START, NURSERY, 3K OR PRE-KINDERGARTEN	KINDERGARTEN through Grade 5	GRADES 6 through 12
Diphtheria and tetanus toxoid-containing vaccine and pertussis vaccine (DTaP/DTp) ²	4 doses	5 doses or 4 doses if the fourth dose was received at age 4 years or older or 3 doses if the child is age 7 years or older and the series was started at age 1 year or older	3 doses
Tetanus and diphtheria toxoid-containing vaccine and pertussis vaccine booster (Tdap) ³	Not Applicable		1 dose
Polio vaccine (IPV/OPV) ^{1,4}	3 doses	4 doses or 3 doses if the third dose was received at age 4 years or older	
Measles, mumps, and rubella vaccine (MMR) ^{1,5}	1 dose	2 doses	
Hepatitis B (HepB) vaccine ^{1,6}	3 doses	3 doses	3 doses or 2 doses of adult hepatitis B vaccine (Recombivax HB [®]) for children who received the doses at least 4 months apart between the ages of 11 through 15 years
Varicella (chickenpox) vaccine ^{1,7}	1 dose	2 doses	
Meningococcal conjugate vaccine (MenACWY) ⁸	Not Applicable		Grades 7, 8, 9, 10 and 11: 1 dose Grade 12: 2 doses or 1 dose if the first dose was received at age 16 years or older
Haemophilus influenzae type b conjugate vaccine (Hib) ⁹	1 to 4 doses	Not Applicable	
Pneumococcal conjugate vaccine (PCV) ¹⁰	1 to 4 doses	Not Applicable	
Influenza ¹¹	1 dose	Not Applicable	

For more information contact:

New York State Department of Health, Bureau of Immunization: 518-473-4437

New York City Department of Health and Mental Hygiene, Bureau of Immunization: 347-396-2433, Office of School Health Citywide (all districts): OSH@health.nyc.gov

- Documented serologic evidence of immunity to measles, mumps, rubella, hepatitis B, or varicella meets the requirements for these immunizations. Serologic evidence of immunity to polio is acceptable only if results are positive for all three serotypes and testing must have been done prior to September 1, 2019. Diagnosis by a physician, physician assistant or nurse practitioner that a child had varicella disease is acceptable proof of immunity to varicella.
- **Diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine – (Minimum age: 6 weeks)**
 - a. Children starting the series on time should receive a five-dose series of DTaP vaccine at ages 2 months, 4 months, 6 months, 15 through 18 months, and age 4 years or older. The fourth dose may be received as early as age 12 months, provided at least 6 months have elapsed since the third dose. However, when retrospectively identified, the fourth dose need not be repeated if it was administered at least 4 months after the third dose. The final dose in the series must be received on or after the fourth birthday and at least 6 months after the prior dose.
 - b. If the fourth dose was administered at age 4 years or older, the fifth (booster) dose is not necessary.
 - c. If the fifth dose was received prior to the fourth birthday, a sixth dose, administered at least 6 months after the prior dose, is required.
 - d. Children ages 7 years and older who are not fully immunized with the childhood DTaP vaccine series should receive Tdap vaccine as the first dose in the catch-up series; if additional doses are needed, either Tdap or Td should be used; the Tdap dose may count towards the Tdap requirement according to grade (see footnote 3d). If the first dose of DTaP/DTp/DT was received before the first birthday, then four total doses are required to complete the series. If the first dose of DTaP/DTp/DT was received on or after the first birthday, then three total doses are required to complete the series. The final dose must be received on or after the fourth birthday.
- **Tetanus and diphtheria toxoids and acellular pertussis (Tdap) vaccine – (Minimum age: 10 years for grades 6-9; 7 years for grades 10-12)**
 - a. Students ages 11 years or older entering grades 6 through 12 are required to have one dose of Tdap.
 - b. Students without Tdap who are age 10 years upon entry to 6th grade are in compliance until they turn age 11 years.
 - c. In addition to the grade 6 through 12 requirement, Tdap may also be given as part of the catch-up series for students 7 years of age and older who are not fully immunized with the childhood DTaP series (see footnote 2e).
 - d. In school year 2023-2024, only doses of Tdap (or DTaP) given at age 10 years or older will satisfy the Tdap requirement for grades 6, 7, 8 and 9. However, doses of Tdap (or DTaP) given at age 7 years or older will satisfy the requirement grades 10 through 12.
 - e. DTaP should NOT be used on or after the 7th birthday but if inadvertently received, the Tdap requirement is satisfied by doses of DTaP (see footnote 3c and 3d).
- **Inactivated poliovirus vaccine (IPV) or oral polio vaccine (OPV) – (Minimum age: 6 weeks)**
 - a. Children starting the series on time should receive IPV at ages 2 months, 4 months, 6 through 18 months and age 4 years or older. The final dose in the series must be received on or after the fourth birthday and at least 6 months after the prior dose.
 - b. For students who received their fourth dose before age 4 years: if the 4th dose was prior to August 7, 2010, four doses separated by at least four weeks is sufficient.
 - c. If the third dose was received at age 4 years or older and at least 6 months after the prior dose, a fourth dose is not necessary.
 - d. If both OPV and IPV were administered as part of a series, the total number of doses and intervals between doses is the same as that recommended for the IPV schedule. For OPV to count towards the completion of the polio series, the dose(s) must have been given before April 1, 2016, and be trivalent (tOPV).
- **Measles, mumps, and rubella (MMR) vaccine – (Minimum age: 12 months)**
 - a. The first dose of MMR vaccine must be given on or after the first birthday. The second dose must be given at least 28 days (four weeks) after the first dose to be considered valid.
 - b. Students in kindergarten through grade 12 must receive two doses of measles-containing vaccine, two doses of mumps-containing vaccine and at least one dose of rubella-containing vaccine.
- **Hepatitis B (HepB) vaccine – (Minimum age: birth)**
 - a. The first dose of HepB vaccine may be given at birth or anytime thereafter. The second dose must be given at least four weeks (28 days) after the first dose. The third dose must be given at least eight weeks after the second dose AND at least 16 weeks after dose one AND no earlier than 24 weeks of age.
 - b. Administration of a total of four doses is permitted when a combination vaccine containing HepB is administered after the birth dose. This fourth dose is often needed to ensure that the last dose in the series is given on or after age 6 months.
 - c. Two doses of adult HepB vaccine (Recombivax[®]) received at least four months apart at age 11 through 15 years will meet the requirement.
- **Varicella (chickenpox) vaccine – (Minimum age: 12 months)**
 - a. The first dose of varicella vaccine must be given on or after the first birthday. The second dose must be given at least 28 days (four weeks) after the first dose to be considered valid.
 - b. For children younger than age 13 years, the recommended minimum interval between doses is three months (though, if the second dose was administered at least four weeks after the first dose, it can be accepted as valid); for people aged 13 years and older, the minimum interval between doses is four weeks.
- **Meningococcal Vaccine (MenACWY) – (Minimum age: 10 years for grades 7-10; 2 months for grades 11 and 12)**
 - a. Students entering grades 7, 8, 9, 10 and 11 are required to receive a single dose of meningococcal conjugate vaccine against serogroups A, C, W-135 and Y (MenACWY vaccines, including Menactra, Merveo, or MenQuadfi). See footnote 8e for the age requirements.
 - b. Students entering grade 12 need to receive two doses of MenACWY vaccine, or only one dose of MenACWY vaccine if the first dose was administered at age 16 years or older.
 - c. If the second dose was administered before age 16 years, then a third dose given on or after age 16 years is required.
 - d. The minimum interval between doses of MenACWY vaccine is eight weeks.
 - e. In school year 2023-2024, only doses of MenACWY given at 10 years or older satisfies the requirement for grades 7, 8, 9 and 10; doses given before 10 years will satisfy the requirement for the first dose for grades 11 and 12.
- **Haemophilus influenzae type b conjugate vaccine (Hib) – (Minimum age: 6 weeks)**
 - a. Children starting the series on time and receiving PRP-T Hib vaccine should receive doses at ages 2 months, 4 months, 6 months and 12 through 15 months. If the formulation is PRP-OMP, only two doses are needed before age 12 through 15 months.
 - b. If 2 doses of vaccine were received before age 12 months, only 3 doses are required, with the third dose at 12 through 15 months and at least 8 weeks after the second dose.
 - c. If the first dose was received at age 12 through 14 months, only 2 doses are required with second dose at least 8 weeks after the first dose.
 - d. If the first dose was received at age 15 months or older, no further doses are required.
 - e. Hib vaccine is not required for children ages 5 years or older.
 - f. For further information, see CDC [catch-up schedule](#)
- **Pneumococcal conjugate vaccine (PCV) – (Minimum age: 6 weeks)**
 - a. Children starting the series on time should receive PCV vaccine at ages 2 months, 4 months, 6 months and 12 through 15 months.
 - b. Unvaccinated children ages 7 through 11 months must receive two doses, at least four weeks apart, followed by a third dose at age 12 through 15 months and at least eight weeks after the prior dose.
 - c. Unvaccinated children ages 12 through 23 months must receive two doses at least eight weeks apart.
 - d. Unvaccinated children ages 24 through 59 months must receive just one dose.
 - e. PCV vaccine is not required for children ages 5 years or older.
 - f. For further information, see CDC [catch-up schedule](#)
- **Influenza Vaccine – (Minimum age: 6 months)**
 - a. All children 6 months through 59 months of age enrolled in NYC Article 47 & 43 regulated Child Care, Head Start, Nursery, or Pre-K programs must receive one dose of influenza vaccine between July 1 and December 31 of each year.
 - b. Depending on their prior influenza vaccination history, some children may need two doses of influenza vaccine; however, a second dose is not required for school entry. Please refer to the Centers for Disease Control and Prevention ([cdc.gov/flu](#)) or New York City Department of Health ([www.nyc.gov/health/flu](#))